

OPHTHALMIC DRUG COMPOSITIONS

Description of corresponding document: **EP0914827**

Technical Field

[0001] The present invention relates to an ophthalmic drug composition containing Phe-Gly-Leu-Met-NH₂ (hereinafter, referred to as FGLM), which is a tetrapeptide at the C-terminal of Substance P, or a pharmaceutically acceptable salt thereof as an active ingredient. More particularly, it relates to a therapeutic agent for corneal disorders having a promoting action on corneal epithelial wound healing, characterized in that an insulin-like growth factor I (hereinafter, referred to as IGF-I), which is one of growth factors, is formulated as another active ingredient with FGLM in a preparation or these two are administered jointly.

Background Art

[0002] Cornea is a transparent, avascular tissue having a diameter of about 1 cm and a thickness of about 1 mm. Transparency of the cornea affects visual functions greatly and various physiological and biochemical phenomena in cornea function mainly with an object of maintenance of transparency of cornea.

[0003] Corneal epithelial defects caused by various diseases such as corneal ulcer, corneal erosion, keratitis and dry eye heals spontaneously unless mixed infection intercurrently occurs. However, if the healing is delayed or does not occur due to some causes and the corneal defects become persistent, the normal construction of the epithelium is badly affected, and further even the structure and function of corneal stroma and endothelium are damaged. The principle of the conventional therapeutic methods is merely passive, i.e., surface of cornea is protected from the outside stimulation whereby epithelium is spontaneously extended to re-cover the defected area. Recent progress of cell biology has revealed factors participating in proliferation, migration, attachment, spreading of cells, etc. and it was reported that compounds which enhance the migration of corneal epithelium play an important role in healing the corneal epithelial defects (Jpn. J. Clin. Ophthalmol., 46, 738-743 (1992); and Jpn. J. Ophthalm. Surg., 5, 719-727 (1992)).

[0004] FGLM is a tetrapeptide at the C-terminal of Substance P which is disclosed in U. S. Patent No. 3,862,114 and the patent specification describes that it has an hypotensive action. Substance P is a polypeptide consisting of eleven amino acids which show actions such as vasodilation, smooth muscle contraction, promotion of salivary gland secretion and diuresis. With regard to Substance P, various studies were conducted in an ophthalmic field. For example, improvement in aberrant conjunctival goblet cell secretion in ophthalmic diseases was disclosed (WO95/13087) and sequential charges of Substance P in inflammation such as keratitis was reported (J. Jpn. Ophthalmol. Soc., 91, 982-987 (1987); *ibid.*, 92, 448-452 (1988)). However, there has been no report on FGLM, which is its partial peptide, in an ophthalmic field.

[0005] On the other hand, an insulin-like growth factor is one of the growth factors controlling the growth of normal human cells such as epidermal growth factor, fibroblast growth factor, platelet-derived growth factor and transforming growth factor, and there are IGF-I and an insulin-like growth factor II (hereinafter, referred to as IGF-II) therein. Recently, it was reported that IGF-I stimulates thyroid cell proliferation (J. Biol. Chem., 264, 18485-18488 (1989)), that IGF-II regulates muscle growth and differentiation (Hum. Mol. Genet., 3, 1117-1121 (1994)), etc. In an ophthalmic field, also, it was disclosed that IGF-I, IGF-II and functional derivatives thereof promote the survival of retinal neurons (cf. Japanese Laid-Open Patent Publication Hei-07/500,839), that IGF-II is effective for therapy of every wound of various areas including the wound upon corneal transplantation (Japanese Laid-Open Patent Publication Sho-63/233,925) and that ocular tissues such as cornea provided to transplantation can be preserved at low temperature in a fresh state of the tissues by using a solution containing the above-mentioned growth factor (Japanese Laid-Open Patent Publications Hei-05/025,001 and Hei-06/048,901). It was also disclosed that, in general, gel compositions containing the growth factor are effective for wound healing in anterior segments etc. (Japanese Laid-Open Patent Publication Hei-02/000,112). However, the growth factors which are specifically disclosed in those patent Publications are epidermal growth factors only and there is no description on the effect of IGF-I.

[0006] IGF-II has been known to be useful for a therapy of wound upon corneal transplantation etc. as mentioned above. However, with regard to IGF-I, there is only a report that it does not affect corneal epithelial wound healing (Connect. Tissue, 27, 65 (1995)).

[0007] It was reported that Substance P itself does not affect the corneal epithelial wound healing but it promotes the corneal epithelial wound healing when it coexists with epidermal growth factor out of growth factor (Prog. Med., 13, 2626-2627 (1993)) or IGF-I (Connect. Tissue, 27, 65 (1995)). However, it has not been revealed that which site of Substance P is an activity-exhibiting site.

[0008] As mentioned above, it has been a very interesting subject to find a minimum activity-exhibiting site of Substance P and to study actions of the compound in such a minimum unit in an ophthalmic field, particularly to study the action on to corneal disorders.

Disclosure of the Invention

[0009] The present inventors have paid their attention to the partial peptide at the C terminal of Substance P and studied its action on corneal disorders. As a result, they have found that FGLM, which is a tetrapeptide at the C terminal of Substance P, promotes the corneal epithelial wound healing when it coexists with IGF-I and that FGLM is a minimum unit of the partial peptide of Substance P for exhibiting the above-mentioned action. Thus, the novel use of FGLM as an ophthalmic drug composition has been found. Further, it has been found that a combined use of FGLM or a pharmaceutically acceptable salt thereof together with IGF-I as another active ingredient is useful for therapy of corneal disorders such as corneal ulcer, corneal erosion, keratitis and dry eye, where cornea is in a wounded state due to various causes.

[0010] FGLM is a tetrapeptide of the C terminal of Substance P and has a structure of Phe-Gly-Leu-Met-NH₂. Phe, Leu and Met have L-, D- and DL-form respectively and all of them belong to the present invention. The preferred embodiment is a compound where all of them are L-form.

[0011] Examples of the pharmaceutically acceptable salt of FGLM are hydrochloride, sulfate, phosphate, lactate, maleate, fumarate, oxalate, methanesulfonate and p-toluenesulfonate.

[0012] The term "corneal disorders" used in the present invention stands for corneal ulcer, corneal erosion, keratitis, dry eye and the like, where the cornea is in a wounded state due to various causes.

[0013] In order to study the usefulness of FGLM and IGF-I, their influence on corneal disorders has been investigated. Although the details will be mentioned later in the item of the "pharmacological tests", it has been found that the coexistence of FGLM and IGF-I promotes the migration of corneal epithelium in a tissue culture system of cornea pieces and the wound healing after corneal erosion. From these findings, it has been revealed that FGLM and IGF-I are useful for the therapy of corneal disorders such as corneal ulcer, corneal erosion, keratitis and dry eye, where the cornea is in a wounded state due to various causes and that they are particularly useful for the therapy of corneal erosion and dry eye.

[0014] FGLM or a pharmaceutically acceptable salt thereof and IGF-I can be administered either orally or parenterally and these two active ingredients can be either formulated together in one preparation or formulated separately, followed by administering them together. Examples of the dosage form are tablets, capsules, granules, powder, injection and ophthalmic preparations and the ophthalmic preparations such as eye drops and eye ointment is particularly preferable. They can be prepared by widely used techniques. In the case of oral preparations such as tablets, capsules, granules and powder, fillers such as lactose, crystalline cellulose, starch and vegetable oil, lubricants such as magnesium stearate and talc, binders such as hydroxypropyl cellulose and polyvinylpyrrolidone, disintegrator such as calcium carboxymethyl cellulose and lowly substituted hydroxypropylmethyl cellulose, coating agents such as hydroxypropylmethyl cellulose, Macrogol and silicone resin, film forming agents such as gelatin and the like can be added thereto, if necessary. In the case of eye drops, isotonic agents such as sodium chloride, buffers such as sodium phosphate, preservatives such as benzalkonium chloride and the like can be used. The pH can be within a range that is acceptable in the ophthalmic preparations and, preferably, it is within a range of 4-8. In the case of eye ointment, widely used bases such as white vaseline and liquid paraffin can be used for the preparation.

[0015] The dose can be appropriately selected depending upon symptom, age, dosage form, etc. Thus, in the case of oral preparations, the doses of FGLM or pharmaceutically acceptable salt thereof and IGF-I

are usually 0.1-5,000 mg (as FGLM) and 0.001-100 mg per day, respectively, preferably 1-1,000mg (as FGLM) and 0.01-10 mg per day, respectively. Administration can be conducted by instilling it once or several times a day. In the case of ophthalmic preparations, concentrations of those active ingredients are 0.001-10% (w/v) (as FGLM) and 0.00001-0.1% (w/v), respectively, preferably 0.01-1% (w/v)(as FGLM) and 0.0001-0.01% (w/v), respectively. Administration can be conducted by instilling it once or several times a day.

[0016] Examples of the preparations and the results of pharmacological tests are given as hereunder although they are intended to give better understanding of the present invention and they do not limit the scope of the present invention.

Best Mode for Carrying Out the Invention

[Examples of the Preparations]

[0017] Representative examples of the preparations used in the present invention will be given hereunder.

1. Ophthalmic Preparations

[0018] The ophthalmic preparations having the following formulations were prepared by widely used methods.

Formulation Example 1. (Eye Drops)

```
<tb><TABLE> Columns=2
<tb>Title: (in 100 ml)
<tb>FGLM<SEP>100 mg
<tb>Sodium chloride<SEP>900 mg
<tb>Sodium hydroxide<SEP>q.s.
<tb>Hydrochloric acid<SEP>q.s.
<tb>Sterile purified water<SEP>q.s.
<tb></TABLE>
```

[0020] Eye drops containing 1 mg, 5 mg, 10 mg, 50 mg, 500 mg and 1,000 mg of FGLM in 100 ml can be prepared as well in the same manner as in Formulation Example 1.

Formulation Example 2. (Eye Drops)

```
<tb><TABLE> Columns=2
<tb>Title: (in 100 ml)
<tb>IGF-I<SEP>1 mg
<tb>Sodium chloride<SEP>900 mg
<tb>Sodium hydroxide<SEP>q.s.
<tb>Hydrochloric acid<SEP>q.s.
<tb>Sterile purified water<SEP>q.s.
<tb></TABLE>
```

[0022] Eye drops containing 0.01 mg, 0.05 mg, 0.1 mg, 0.5 mg, 5 mg, 10 mg, 50 mg and 100 mg of IGF-I in 100 ml can be prepared as well in the same manner as in Formulation Example 2.

Formulation Example 3 (Eye Drops)

```
<tb><TABLE> Columns=2
<tb>Title: (in 100 ml)
```

<tb>FGLM<SEP>100 mg
<tb>IGF-I<SEP>1 mg
<tb>Sodium chloride<SEP>900 mg
<tb>Sodium hydroxide<SEP>q.s.
<tb>Hydrochloric acid<SEP>q.s.
<tb>Sterile purified water<SEP>q.s.
<tb></TABLE>

[0024] Eye drops containing 1 mg, 5 mg, 10 mg, 50 mg, 500 mg and 1,000 mg of FGLM and 0.01 mg, 0.05 mg, 0.1 mg, 0.5 mg, 10 mg, 50 mg and 100 mg of IGF-I in any combination thereof can be prepared as well in the same manner as in Formulation Example 3.

Formulation Example 4. (Eye Ointment)

<tb><TABLE> Columns=2
<tb>Title: (in 100 g)
<tb>FGLM<SEP>100 mg
<tb>IGF-I<SEP>1 mg
<tb>White vaseline<SEP>90 g
<tb>Liquid paraffin<SEP>q.s
<tb></TABLE>

[0026] Eye ointments containing 1 mg, 5 mg, 10 mg, 50 mg, 100 mg, 500 mg and 1,000 mg of FGLM and 0.01 mg, 0.05 mg, 0.1 mg, 0.5 mg, 1 mg, 10 mg, 50 mg and 100 mg of IGF-I in any combination thereof can be prepared as well in the same manner as in Formulation Example 4,.

2. Tablets

[0027] Tablets of the following formulations were prepared by widely used methods.

Formulation Example 5.

<tb><TABLE> Columns=2
<tb>Title: (in 100 mg)
<tb>FGLM<SEP>10 mg
<tb>Lactose<SEP>59.4 mg
<tb>Corn starch<SEP>20 mg
<tb>Calcium carboxymethyl cellulose<SEP>6 mg
<tb>Hydroxypropyl cellulose<SEP>4 mg
<tb>Magnesium stearate<SEP>0.6 mg
<tb></TABLE>

[0029] The tablet of the above formulation can be coated with 2 mg of a coating agent such as hydroxypropyl cellulose.

[0030] Tablets containing 0.1 mg, 0.5 mg, 1 mg, 5 mg and 50 mg of FGLM in 100 mg of a tablet can be prepared as well in the same manner as in Formulation Example 5.

Formulation Example 6.

<tb><TABLE> Columns=2
<tb>Title: (in 100 mg)
<tb>IGF-I<SEP>0.1 mg
<tb>Lactose<SEP>69.3 mg
<tb>Corn starch<SEP>20 mg
<tb>Calcium carboxymethyl cellulose<SEP>6 mg
<tb>Hydroxypropyl cellulose<SEP>4 mg
<tb>Magnesium stearate<SEP>0.6 mg
<tb></TABLE>

[0032] The tablet of the above formulation can be coated with 2 mg of a coating agent such as hydroxypropyl cellulose.

[0033] Tablets containing 0.001 mg, 0.01 mg, 0.05 mg, 0.5 mg, 1 mg, 5 mg, 10 mg and 50 mg of IGF-I in 100 mg of a tablet can be prepared as well in the same manner as in Formulation Example 6.

Formulation Example 7.

```
<tb><TABLE> Columns=2
<tb>Title: (in 100 mg)
<tb>FGLM<SEP>10 mg
<tb>IGF-I<SEP>0.1 mg
<tb>Lactose<SEP>59.3 mg
<tb>Corn starch<SEP>20 mg
<tb>Calcium carboxymethyl cellulose<SEP>6 mg
<tb>Hydroxypropyl cellulose<SEP>4 mg
<tb>Magnesium stearate<SEP>0.6 mg
<tb></TABLE>
```

[0035] The tablet of the above formulation can be coated with 2 mg of a coating agent such as hydroxypropyl cellulose.

[0036] Tablets containing 0.1 mg, 0.5 mg, 1 mg, 5 mg and 10 mg of FGLM and 0.001 mg, 0.01 mg, 0.05 mg, 0.1 mg, 0.5 mg, 1 mg, 5 mg and 10 mg of IGF-I in 100 mg of a tablet in any combination thereof can be prepared as well in the same manner as in Formulation Example 7.

[Pharmacological Tests]

1) Corneal Epithelial Migration (in vitro)

[0037] Cornea of male Japanese white rabbits was used. The influence on migration of corneal epithelium was investigated using as an index the extended length of cornea epithelium in a tissue culture system of the cornea pieces according to a method of Nishida et al (J. Cell Biol., 97, 1653-1657 (1983)).

(Experimental Method)

[0038] Cornea blocks (six blocks per group) cut out from cornea pieces of the rabbit were incubated in a culture medium (TC-199) containing the test compound under the condition of 37 DEG C and 5% of CO₂ for 24 hours. After the incubation, the cornea blocks were fixed in a mixed solution of ethanol and glacial acetic acid (95:5 by volume) and embedded in paraffin to prepare the slices. Paraffin was removed from the slices, then the slices were stained with hematoxyline and eosin and the extended length of the epithelial cell layer was measured under a microscope.

[0039] The product which was prepared by the same incubation using a culture medium containing no test compound was used as a control.

(Results)

[0040] As an example of the results of the experiment, Table 1 shows the result where incubation was conducted using a culture medium containing FGLM alone, IGF-I alone and both FGLM and IGF-I. Table 2 shows the result where the peptide which was added to the culture medium was Gly-Leu-Met-NH₂ (hereinafter, referred to as GLM), FGLM, Val-Gly-Leu-Met-NH₂ (hereinafter, referred to as VGLM), Ile-Gly-Leu-Met-NH₂ (hereinafter, referred to as IGLM), Tyr-Gly-Leu-Met-NH₂ (hereinafter, referred to as YGLM) or Phe-Phe-Gly-Leu-Met-NH₂ (hereinafter, referred to as FFGLM).

```

<tb><TABLE> Id=Table 1 Columns=2
<tb>
<tb>Head Col 1:
<tb>Head Col 2: Extended Length (  $\mu$  m)
<tb>Control<SEP>433
<tb>FGLM (20  $\mu$  M)<SEP>426
<tb>IGF-I (10 ng/ml)<SEP>430
<tb>FGLM (20  $\mu$  M) and IGF-I (10 ng/ml)<SEP>662
<tb></TABLE>
<tb><TABLE> Id=Table 2 Columns=2
<tb>
<tb>Head Col 1:
<tb>Head Col 2: Extended Length (  $\mu$  m)
<tb>Control<SEP>433
<tb>IGF-I (10 ng/ml)<SEP>430
<tb>IGF-I (10 ng/ml) and GLM (20  $\mu$  M)<SEP>445
<tb>IGF-I (10 ng/ml) and FGLM (20  $\mu$  M)<SEP>662
<tb>IGF-I (10 ng/ml) and VGLM (20  $\mu$  M)<SEP>440
<tb>IGF-I (10 ng/ml) and IGLM (20  $\mu$  M)<SEP>426
<tb>IGF-I (10 ng/ml) and YGLM (20  $\mu$  M)<SEP>433
<tb>IGF-I (10 ng/ml) and FFGLM (20  $\mu$  M)<SEP>655
<tb></TABLE>

```

[0041] As shown in Table 1, while no influence on migration of corneal epithelium was noted by a sole administration of FGLM or of IGF-I, when incubation was conducted in a culture medium containing both FGLM and IGF-I, a remarkable promotion of the migration of corneal epithelium was noted.

[0042] Further, as shown in Table 2, with regard to the peptide which was added to the culture medium together with IGF-I, while a remarkable promotion of the migration of corneal epithelium was noted when FGLM or FFGLM was used, no influence on the migration of corneal epithelium was noted when the C-terminal tripeptide of Substance P or a peptide similar to FGLM was added.

2) Promoting Action on Corneal Wound Healing (in vivo)

[0043] Male Japanese white rabbits were used. Corneal erosion was caused according to a method of Cintron et al (Ophthalmic Res., 11, 90-96 (1979)). The wounded area was measured using an area stained with fluorescein as an index and the influence on the corneal wound healing was investigated.

(Experimental Method)

[0044] After the onset of corneal erosion, eye drops containing the test compound were instilled every two hours six times a day (50 μ l per administration). In measuring the wounded area, a fluorescein staining was conducted and photographs of the cornea were taken and measured. Area of the cornea stained with fluorescein in the photographs was calculated by means of an image analysis processing system.

[0045] Rabbit to which a base containing no test compound was instilled was used as a control.

(Results)

[0046] As an example of the results of the experiment, Table 3 shows wounded areas of immediately after onset of corneal erosion and 12, 24, 36 and 48 hours thereafter upon the treatment where eye drops containing 0.05% (w/v) of FGLM (F-3) alone was instilled; where eye drops containing 0.0001% (w/v) of IGF-I (I-6) alone was instilled; and where both eye drops containing 0.05% (w/v) of FGLM (F-3) and that containing 0.0001% (w/v) of IGF-I (I-6) were instilled.

```

<tb><TABLE> Id=Table 3 Columns=6
<tb>
<tb>Head Col 1:
<tb>Head Col 2 to 6: Wounded Area (mm<sup>2</sup>) after Corneal Erosion

```

<tb>
<tb>SubHead Col 1:
<tb>SubHead Col 2: 0 hr
<tb>SubHead Col 3: 12 hrs
<tb>SubHead Col 4: 24 hrs
<tb>SubHead Col 5: 36 hrs
<tb>SubHead Col 6: 48 hrs
<tb>Control<SEP>35.4<SEP>31.6<SEP>20.6<SEP>11.7<SEP>3.3
<tb>FGLM<SEP>35.4<SEP>31.0<SEP>19.9<SEP>10.9<SEP>2.9
<tb>IGF-I<SEP>35.5<SEP>30.3<SEP>19.0<SEP>10.0<SEP>2.6
<tb>FGLM + IGF-I<SEP>35.5<SEP>28.1<SEP>10.5<SEP>2.4<SEP>0.1
<tb></TABLE>

[0047] As shown in Table 3, while no influence was noted on wound healing after corneal erosion by a sole administration of FGLM or IGF-I, when both FGLM and IGF-I were instilled, a remarkable promotion was noted in wound healing.

[0048] From the above-mentioned pharmacological tests, it has been found that, when FGLM, which is one of the partial peptides of Substance P, or a pharmaceutically acceptable salt thereof coexists with IGF-I, which is one of the growth factors, the former exhibits an promoting action on the corneal epithelial wound healing whereby it is useful as a therapeutic agent for corneal disorders where cornea is wounded due to various causes such as corneal ulcer, corneal erosion, keratitis and dry eye.

[0049] In addition, while the promoting action on the corneal epithelial migration was noted in tetrapeptide or pentapeptide at the C-terminal of Substance P when it coexists with IGF-I, this action was not noted in tripeptide at the C-terminal of Substance P. Accordingly, it has been now revealed that the minimum unit of the partial peptide of Substance P coexisting with IGF-I, which is necessary for exhibiting the promoting action on corneal epithelial wound healing, was the tetrapeptide at the C-terminal. Further, when the amino acid at the N-terminal of the tetrapeptide was other than Phe, no promoting action on the corneal epithelial migration was noted. Accordingly, it has been revealed that the tetrapeptide coexisting with IGF-I which is necessary for exhibiting the promoting action on corneal epithelial wound healing should be FGLM, which is a tetrapeptide at the C-terminal of Substance P.

Data supplied from the **esp@cenet** database - Worldwide

OPHTHALMIC DRUG COMPOSITIONS

Claims of corresponding document: **EP0914827**

1. An ophthalmic composition comprising Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof as an active ingredient.
2. A therapeutic agent for corneal disorder comprising Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof as an active ingredient.
3. A therapeutic agent for corneal disorder comprising Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof and an insulin-like growth factor I as active ingredients.
4. The therapeutic agent for corneal disorder according to claim 2 or 3 where the corneal disorder is corneal ulcer, corneal erosion, keratitis or dry eye.
5. the therapeutic agent for corneal disorder according to claim 2 or 3 where the corneal disorder is corneal erosion or dry eye.
6. The therapeutic agent for corneal disorder according to any of claims 2 to 5 where the dosage form is an ophthalmic preparation.
7. A promoting agent for corneal epithelial migration comprising Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof and an insulin-like growth factor I as active ingredients.
8. The promoting agent for corneal epithelial migration according to claim 7 where the dosage form is an ophthalmic preparation.

Data supplied from the **esp@cenet** database - Worldwide

AI

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 914 827 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

12.05.1999 Bulletin 1999/19

(51) Int. Cl.⁶: **A61K 38/07**

(21) Application number: 97926223.5

(86) International application number:

PCT/JP97/02015

(22) Date of filing: 11.06.1997

(87) International publication number:

WO 97/49419 (31.12.1997 Gazette 1997/57)

(84) Designated Contracting States:

AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

(30) Priority: 26.06.1996 JP 165612/96

(71) Applicants:

- SANTEN PHARMACEUTICAL CO., LTD.
Higashiyodogawa-ku, Osaka-shi, Osaka 533-
8651 (JP)
- Nishida, Teruo
Ube-shi, Yamaguchi 755 (JP)

(72) Inventors:

- NISHIDA, Teruo
Ube-shi, Yamaguchi 755 (JP)
- NAKAMURA, Masatsugu
Nara-shi, Nara 631 (JP)
- NAKATA, Katsuhiko
Sakurai-shi, Nara 633 (JP)

(74) Representative:

Barske, Helko, Dr. rer. nat.
Blumbach, Kramer & Partner
Patentanwälte
Radeckestrasse 43
81245 München (DE)

(54) OPTHALMIC DRUG COMPOSITIONS

(57) An object of the present invention is to find a minimum activity exhibiting sites of Substance P, to reveal the action of the compound of the minimum unit in an ophthalmic field and to offer an ophthalmic composition comprising the compound as an active ingredient. The present invention relates to an ophthalmic composition, particularly, to a therapeutic agent for corneal disorder which comprises Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof as an active ingredient. The present invention also relates to a therapeutic agent for corneal disorder, particularly, to a promoting agent for corneal epithelial migration which comprises Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof and an insulin-like growth factor I as active ingredients. Preferred dosage form is an ophthalmic preparation.

EP 0 914 827 A1

Description

Technical Field

5 [0001] The present invention relates to an ophthalmic drug composition containing Phe-Gly-Leu-Met-NH₂ (hereinafter, referred to as FGLM), which is a tetrapeptide at the C-terminal of Substance P, or a pharmaceutically acceptable salt thereof as an active ingredient. More particularly, it relates to a therapeutic agent for corneal disorders having a promoting action on corneal epithelial wound healing, characterized in that an insulin-like growth factor I (hereinafter, referred to as IGF-I), which is one of growth factors, is formulated as another active ingredient with FGLM in a preparation or these two are administered jointly.

Background Art

15 [0002] Cornea is a transparent, avascular tissue having a diameter of about 1 cm and a thickness of about 1 mm. Transparency of the cornea affects visual functions greatly and various physiological and biochemical phenomena in cornea function mainly with an object of maintenance of transparency of cornea.

[0003] Corneal epithelial defects caused by various diseases such as corneal ulcer, corneal erosion, keratitis and dry eye heals spontaneously unless mixed infection intercurrently occurs. However, if the healing is delayed or does not occur due to some causes and the corneal defects become persistent, the normal construction of the epithelium is badly affected, and further even the structure and function of corneal stroma and endothelium are damaged. The principle of the conventional therapeutic methods is merely passive, i.e., surface of cornea is protected from the outside stimulation whereby epithelium is spontaneously extended to re-cover the defected area. Recent progress of cell biology has revealed factors participating in proliferation, migration, attachment, spreading of cells, etc. and it was reported that compounds which enhance the migration of corneal epithelium play an important role in healing the corneal epithelial defects (*Jpn. J. Clin. Ophthalmol.*, 46, 738-743 (1992); and *Jpn. J. Ophthalm. Surg.*, 5, 719-727 (1992)).

25 [0004] FGLM is a tetrapeptide at the C-terminal of Substance P which is disclosed in U. S. Patent No. 3,862,114 and the patent specification describes that it has an hypotensive action. Substance P is a polypeptide consisting of eleven amino acids which show actions such as vasodilation, smooth muscle contraction, promotion of salivary gland secretion and diuresis. With regard to Substance P, various studies were conducted in an ophthalmic field. For example, improvement in aberrant conjunctival goblet cell secretion in ophthalmic diseases was disclosed (WO95/13087) and sequential charges of Substance P in inflammation such as keratitis was reported (*J. Jpn. Ophthalmol. Soc.*, 91, 982-987 (1987); *ibid.*, 92, 448-452 (1988)). However, there has been no report on FGLM, which is its partial peptide, in an ophthalmic field.

35 [0005] On the other hand, an insulin-like growth factor is one of the growth factors controlling the growth of normal human cells such as epidermal growth factor, fibroblast growth factor, platelet-derived growth factor and transforming growth factor, and there are IGF-I and an insulin-like growth factor II (hereinafter, referred to as IGF-II) therein. Recently, it was reported that IGF-I stimulates thyroid cell proliferation (*J. Biol. Chem.*, 264, 18485-18488 (1989)), that IGF-II regulates muscle growth and differentiation (*Hum. Mol. Genet.*, 3, 1117-1121 (1994)), etc. In an ophthalmic field, also, it was disclosed that IGF-I, IGF-II and functional derivatives thereof promote the survival of retinal neurons (cf. Japanese Laid-Open Patent Publication Hei-07/500,839), that IGF-II is effective for therapy of every wound of various areas including the wound upon corneal transplantation (Japanese Laid-Open Patent Publication Sho-63/233,925) and that ocular tissues such as cornea provided to transplantation can be preserved at low temperature in a fresh state of the tissues by using a solution containing the above-mentioned growth factor (Japanese Laid-Open Patent Publications Hei-05/025,001 and Hei-06/048,901). It was also disclosed that, in general, gel compositions containing the growth factor are effective for wound healing in anterior segments etc. (Japanese Laid-Open Patent Publication Hei-02/000,112). However, the growth factors which are specifically disclosed in those patent Publications are epidermal growth factors only and there is no description on the effect of IGF-I.

45 [0006] IGF-II has been known to be useful for a therapy of wound upon corneal transplantation etc. as mentioned above. However, with regard to IGF-I, there is only a report that it does not affect corneal epithelial wound healing (*Connect. Tissue*, 27, 65 (1995)).

50 [0007] It was reported that Substance P itself does not affect the corneal epithelial wound healing but it promotes the corneal epithelial wound healing when it coexists with epidermal growth factor or growth factor (*Prog. Med.*, 13, 2626-2627 (1993)) or IGF-I (*Connect. Tissue*, 27, 65 (1995)). However, it has not been revealed that which site of Substance P is an activity-exhibiting site.

55 [0008] As mentioned above, it has been a very interesting subject to find a minimum activity-exhibiting site of Substance P and to study actions of the compound in such a minimum unit in an ophthalmic field, particularly to study the action on to corneal disorders.

Disclosure of the Invention

[0009] The present inventors have paid their attention to the partial peptide at the C terminal of Substance P and studied its action on corneal disorders. As a result, they have found that FGLM, which is a tetrapeptide at the C terminal of Substance P, promotes the corneal epithelial wound healing when it coexists with IGF-I and that FGLM is a minimum unit of the partial peptide of Substance P for exhibiting the above-mentioned action. Thus, the novel use of FGLM as an ophthalmic drug composition has been found. Further, it has been found that a combined use of FGLM or a pharmaceutically acceptable salt thereof together with IGF-I as another active ingredient is useful for therapy of corneal disorders such as corneal ulcer, corneal erosion, keratitis and dry eye, where cornea is in a wounded state due to various causes.

[0010] FGLM is a tetrapeptide of the C terminal of Substance P and has a structure of Phe-Gly-Leu-Met-NH₂. Phe, Leu and Met have L-, D- and DL-form respectively and all of them belong to the present invention. The preferred embodiment is a compound where all of them are L-form.

[0011] Examples of the pharmaceutically acceptable salt of FGLM are hydrochloride, sulfate, phosphate, lactate, maleate, fumarate, oxalate, methanesulfonate and p-toluenesulfonate.

[0012] The term "corneal disorders" used in the present invention stands for corneal ulcer, corneal erosion, keratitis, dry eye and the like, where the cornea is in a wounded state due to various causes.

[0013] In order to study the usefulness of FGLM and IGF-I, their influence on corneal disorders has been investigated. Although the details will be mentioned later in the item of the "pharmacological tests", it has been found that the coexistence of FGLM and IGF-I promotes the migration of corneal epithelium in a tissue culture system of cornea pieces and the wound healing after corneal erosion. From these findings, it has been revealed that FGLM and IGF-I are useful for the therapy of corneal disorders such as corneal ulcer, corneal erosion, keratitis and dry eye, where the cornea is in a wounded state due to various causes and that they are particularly useful for the therapy of corneal erosion and dry eye.

[0014] FGLM or a pharmaceutically acceptable salt thereof and IGF-I can be administered either orally or parenterally and these two active ingredients can be either formulated together in one preparation or formulated separately, followed by administering them together. Examples of the dosage form are tablets, capsules, granules, powder, injection and ophthalmic preparations and the ophthalmic preparations such as eye drops and eye ointment is particularly preferable. They can be prepared by widely used techniques. In the case of oral preparations such as tablets, capsules, granules and powder, fillers such as lactose, crystalline cellulose, starch and vegetable oil, lubricants such as magnesium stearate and talc, binders such as hydroxypropyl cellulose and polyvinylpyrrolidone, disintegrator such as calcium carboxymethyl cellulose and lowly substituted hydroxypropylmethyl cellulose, coating agents such as hydroxypropylmethyl cellulose, Macrogol and silicone resin, film forming agents such as gelatin and the like can be added thereto, if necessary. In the case of eye drops, isotonic agents such as sodium chloride, buffers such as sodium phosphate, preservatives such as benzalkonium chloride and the like can be used. The pH can be within a range that is acceptable in the ophthalmic preparations and, preferably, it is within a range of 4-8. In the case of eye ointment, widely used bases such as white vaseline and liquid paraffin can be used for the preparation.

[0015] The dose can be appropriately selected depending upon symptom, age, dosage form, etc. Thus, in the case of oral preparations, the doses of FGLM or pharmaceutically acceptable salt thereof and IGF-I are usually 0.1-5,000 mg (as FGLM) and 0.001-100 mg per day, respectively, preferably 1-1,000mg (as FGLM) and 0.01-10 mg per day, respectively. Administration can be conducted by instilling it once or several times a day. In the case of ophthalmic preparations, concentrations of those active ingredients are 0.001-10% (w/v) (as FGLM) and 0.00001-0.1% (w/v), respectively, preferably 0.01-1% (w/v) (as FGLM) and 0.0001-0.01% (w/v), respectively. Administration can be conducted by instilling it once or several times a day.

[0016] Examples of the preparations and the results of pharmacological tests are given as hereunder although they are intended to give better understanding of the present invention and they do not limit the scope of the present invention.

Best Mode for Carrying Out the Invention

[Examples of the Preparations]

[0017] Representative examples of the preparations used in the present invention will be given hereunder.

1. Ophthalmic Preparations

[0018] The ophthalmic preparations having the following formulations were prepared by widely used methods.

EP 0 914 827 A1

Formulation Example 1. (Eye Drops)

[0019]

(in 100 ml)	
FGLM	100 mg
Sodium chloride	900 mg
Sodium hydroxide	q.s.
Hydrochloric acid	q.s.
Sterile purified water	q.s.

[0020] Eye drops containing 1 mg, 5 mg, 10 mg, 50 mg, 500 mg and 1,000 mg of FGLM in 100 ml can be prepared as well in the same manner as in Formulation Example 1.

Formulation Example 2. (Eye Drops)

[0021]

(in 100 ml)	
IGF-I	1 mg
Sodium chloride	900 mg
Sodium hydroxide	q.s.
Hydrochloric acid	q.s.
Sterile purified water	q.s.

[0022] Eye drops containing 0.01 mg, 0.05 mg, 0.1 mg, 0.5 mg, 5 mg, 10 mg, 50 mg and 100 mg of IGF-I in 100 ml can be prepared as well in the same manner as in Formulation Example 2.

Formulation Example 3 (Eye Drops)

[0023]

(in 100 ml)	
FGLM	100 mg
IGF-I	1 mg
Sodium chloride	900 mg
Sodium hydroxide	q.s.
Hydrochloric acid	q.s.
Sterile purified water	q.s.

[0024] Eye drops containing 1 mg, 5 mg, 10 mg, 50 mg, 500 mg and 1,000 mg of FGLM and 0.01 mg, 0.05 mg, 0.1 mg, 0.5 mg, 10 mg, 50 mg and 100 mg of IGF-I in any combination thereof can be prepared as well in the same manner

EP 0 914 827 A1

as in Formulation Example 3.

Formulation Example 4. (Eye Ointment)

[0025]

(in 100 g)	
FGLM	100 mg
IGF-I	1 mg
White vaseline	90 g
Liquid paraffin	q.s

[0026] Eye ointments containing 1 mg, 5 mg, 10 mg, 50 mg, 100 mg, 500 mg and 1,000 mg of FGLM and 0.01 mg, 0.05 mg, 0.1 mg, 0.5 mg, 1 mg, 10 mg, 50 mg and 100 mg of IGF-I in any combination thereof can be prepared as well in the same manner as in Formulation Example 4,.

2. Tablets

[0027] Tablets of the following formulations were prepared by widely used methods.

Formulation Example 5.

[0028]

(in 100 mg)	
FGLM	10 mg
Lactose	59.4 mg
Corn starch	20 mg
Calcium carboxymethyl cellulose	6 mg
Hydroxypropyl cellulose	4 mg
Magnesium stearate	0.6 mg

[0029] The tablet of the above formulation can be coated with 2 mg of a coating agent such as hydroxypropyl cellulose.

[0030] Tablets containing 0.1 mg, 0.5 mg, 1 mg, 5 mg and 50 mg of FGLM in 100 mg of a tablet can be prepared as well in the same manner as in Formulation Example 5.

Formulation Example 6.

[0031]

(in 100 mg)	
IGF-I	0.1 mg

EP 0 914 827 A1

(continued)

(in 100 mg)	
Lactose	69.3 mg
Corn starch	20 mg
Calcium carboxymethyl cellulose	6 mg
Hydroxypropyl cellulose	4 mg
Magnesium stearate	0.6 mg

[0032] The tablet of the above formulation can be coated with 2 mg of a coating agent such as hydroxypropyl cellulose.

[0033] Tablets containing 0.001 mg, 0.01 mg, 0.05 mg, 0.5 mg, 1 mg, 5 mg, 10 mg and 50 mg of IGF-I in 100 mg of a tablet can be prepared as well in the same manner as in Formulation Example 6.

Formulation Example 7.

[0034]

(in 100 mg)	
FGLM	10 mg
IGF-I	0.1 mg
Lactose	59.3 mg
Corn starch	20 mg
Calcium carboxymethyl cellulose	6 mg
Hydroxypropyl cellulose	4 mg
Magnesium stearate	0.6 mg

[0035] The tablet of the above formulation can be coated with 2 mg of a coating agent such as hydroxypropyl cellulose.

[0036] Tablets containing 0.1 mg, 0.5 mg, 1 mg, 5 mg and 10 mg of FGLM and 0.001 mg, 0.01 mg, 0.05 mg, 0.1 mg, 0.5 mg, 1 mg, 5 mg and 10 mg of IGF-I in 100 mg of a tablet in any combination thereof can be prepared as well in the same manner as in Formulation Example 7.

[Pharmacological Tests]

1) Corneal Epithelial Migration (*in vitro*)

[0037] Cornea of male Japanese white rabbits was used. The influence on migration of corneal epithelium was investigated using as an index the extended length of cornea epithelium in a tissue culture system of the cornea pieces according to a method of Nishida et al (*J. Cell Biol.*, 97, 1653-1657 (1983)).

(Experimental Method)

[0038] Cornea blocks (six blocks per group) cut out from cornea pieces of the rabbit were incubated in a culture medium (TC-199) containing the test compound under the condition of 37°C and 5% of CO₂ for 24 hours. After the incubation, the cornea blocks were fixed in a mixed solution of ethanol and glacial acetic acid (95:5 by volume) and embedded in paraffin to prepare the slices. Paraffin was removed from the slices, then the slices were stained with hematoxyline and eosin and the extended length of the epithelial cell layer was measured under a microscope.

[0039] The product which was prepared by the same incubation using a culture medium containing no test compound was used as a control.

(Results)

[0040] As an example of the results of the experiment, Table 1 shows the result where incubation was conducted using a culture medium containing FGLM alone, IGF-I alone and both FGLM and IGF-I. Table 2 shows the result where the peptide which was added to the culture medium was Gly-Leu-Met-NH₂ (hereinafter, referred to as GLM), FGLM, Val-Gly-Leu-Met-NH₂ (hereinafter, referred to as VGLM), Ile-Gly-Leu-Met-NH₂ (hereinafter, referred to as IGLM), Tyr-Gly-Leu-Met-NH₂ (hereinafter, referred to as YGLM) or Phe-Phe-Gly-Leu-Met-NH₂ (hereinafter, referred to as FFGLM).

Table 1

	Extended Length (μm)
Control	433
FGLM (20 μM)	426
IGF-I (10 ng/ml)	430
FGLM (20 μM) and IGF-I (10 ng/ml)	662

Table 2

	Extended Length (μm)
Control	433
IGF-I (10 ng/ml)	430
IGF-I (10 ng/ml) and GLM (20 μM)	445
IGF-I (10 ng/ml) and FGLM (20 μM)	662
IGF-I (10 ng/ml) and VGLM (20 μM)	440
IGF-I (10 ng/ml) and IGLM (20 μM)	426
IGF-I (10 ng/ml) and YGLM (20 μM)	433
IGF-I (10 ng/ml) and FFGLM (20 μM)	655

[0041] As shown in Table 1, while no influence on migration of corneal epithelium was noted by a sole administration of FGLM or of IGF-I, when incubation was conducted in a culture medium containing both FGLM and IGF-I, a remarkable promotion of the migration of corneal epithelium was noted.

[0042] Further, as shown in Table 2, with regard to the peptide which was added to the culture medium together with IGF-I, while a remarkable promotion of the migration of corneal epithelium was noted when FGLM or FFGLM was used, no influence on the migration of corneal epithelium was noted when the C-terminal tripeptide of Substance P or a peptide similar to FGLM was added.

2) Promoting Action on Corneal Wound Healing (*in vivo*)

[0043] Male Japanese white rabbits were used. Corneal erosion was caused according to a method of Cintron et al (*Ophthalmic Res.*, **11**, 90-96 (1979)). The wounded area was measured using an area stained with fluorescein as an index and the influence on the corneal wound healing was investigated.

(Experimental Method)

[0044] After the onset of corneal erosion, eye drops containing the test compound were instilled every two hours six times a day (50 μl per administration). In measuring the wounded area, a fluorescein staining was conducted and photographs of the cornea were taken and measured. Area of the cornea stained with fluorescein in the photographs was calculated by means of an image analysis processing system.

[0045] Rabbit to which a base containing no test compound was instilled was used as a control.

(Results)

[0046] As an example of the results of the experiment, Table 3 shows wounded areas of immediately after onset of corneal erosion and 12, 24, 36 and 48 hours thereafter upon the treatment where eye drops containing 0.05% (w/v) of FGLM (F-3) alone was instilled; where eye drops containing 0.0001% (w/v) of IGF-I (I-6) alone was instilled; and where both eye drops containing 0.05% (w/v) of FGLM (F-3) and that containing 0.0001% (w/v) of IGF-I (I-6) were instilled.

Table 3

	Wounded Area (mm ²) after Corneal Erosion				
	0 hr	12 hrs	24 hrs	36 hrs	48 hrs
Control	35.4	31.6	20.6	11.7	3.3
FGLM	35.4	31.0	19.9	10.9	2.9
IGF-I	35.5	30.3	19.0	10.0	2.6
FGLM + IGF-I	35.5	28.1	10.5	2.4	0.1

[0047] As shown in Table 3, while no influence was noted on wound healing after corneal erosion by a sole administration of FGLM or IGF-I, when both FGLM and IGF-I were instilled, a remarkable promotion was noted in wound healing.

[0048] From the above-mentioned pharmacological tests, it has been found that, when FGLM, which is one of the partial peptides of Substance P, or a pharmaceutically acceptable salt thereof coexists with IGF-I, which is one of the growth factors, the former exhibits an promoting action on the corneal epithelial wound healing whereby it is useful as a therapeutic agent for corneal disorders where cornea is wounded due to various causes such as corneal ulcer, corneal erosion, keratitis and dry eye.

[0049] In addition, while the promoting action on the corneal epithelial migration was noted in tetrapeptide or pentapeptide at the C-terminal of Substance P when it coexists with IGF-I, this action was not noted in tripeptide at the C-terminal of Substance P. Accordingly, it has been now revealed that the minimum unit of the partial peptide of Substance P coexisting with IGF-I, which is necessary for exhibiting the promoting action on corneal epithelial wound healing, was the tetrapeptide at the C-terminal. Further, when the amino acid at the N-terminal of the tetrapeptide was other than Phe, no promoting action on the corneal epithelial migration was noted. Accordingly, it has been revealed that the tetrapeptide coexisting with IGF-I which is necessary for exhibiting the promoting action on corneal epithelial wound healing should be FGLM, which is a tetrapeptide at the C-terminal of Substance P.

Claims

1. An ophthalmic composition comprising Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof as an active ingredient.
2. A therapeutic agent for corneal disorder comprising Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof as an active ingredient.
3. A therapeutic agent for corneal disorder comprising Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof and an insulin-like growth factor I as active ingredients.
4. The therapeutic agent for corneal disorder according to claim 2 or 3 where the corneal disorder is corneal ulcer, corneal erosion, keratitis or dry eye.
5. the therapeutic agent for corneal disorder according to claim 2 or 3 where the corneal disorder is corneal erosion or dry eye.
6. The therapeutic agent for corneal disorder according to any of claims 2 to 5 where the dosage form is an ophthalmic preparation.

EP 0 914 827 A1

7. A promoting agent for corneal epithelial migration comprising Phe-Gly-Leu-Met-NH₂ or a pharmaceutically acceptable salt thereof and an insulin-like growth factor I as active ingredients.

5 8. The promoting agent for corneal epithelial migration according to claim 7 where the dosage form is an ophthalmic preparation.

10

15

20

25

30

35

40

45

50

55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP97/02015

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl⁶ A61K38/07

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int. Cl⁶ A61K38/07

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Connect. Tissue, Vol. 27, No. 1 (1995), p. 65	1 - 8
A	US, 3862114, A (ICI Australia Ltd.), January 21, 1975 (21. 01. 75) (Family: none)	1 - 8
PA	US, 5616562, A (Murphy, C.J. and others), April 1, 1997 (01. 04. 97) (Family: none) (Refer to column 3, lines 25 to 41; column 5, line 22 to column 6, line 7; example 1; Figs. 1 to 3)	1 - 8

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

September 1, 1997 (01. 09. 97)

Date of mailing of the international search report

September 9, 1997 (09. 09. 97)

Name and mailing address of the ISA/

Japanese Patent Office

Facsimile No.

Authorized officer

Telephone No.